



## Genetic diversity of Treponemes in dairy herds and their surrounding environment

Weiss Nielsen , Martin ; Jensen, Tim Kåre; Ingerslev, Hans-Christian; Schou, Kirstine Klitgaard

*Publication date:*  
2014

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Weiss Nielsen , M., Jensen, T. K., Ingerslev, H-C., & Schou, K. K. (2014). *Genetic diversity of Treponemes in dairy herds and their surrounding environment*. Poster session presented at Gordon Research Conference on Spirochetes, Ventura, CA, United States.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# Genetic diversity of *Treponemes* in dairy herds and their surrounding environment



Martin W. Nielsen, Tim K. Jensen, Hans-Christian Ingerslev, Kirstine Klitgaard

National Veterinary Institute, Section for Bacteriology, Pathology and Parasitology, The Technical University of Denmark

## Introduction

The causative agents of the highly contagious inflammatory skin disease digital dermatitis (DD), which has reached endemic levels in parts of the world, have been identified as mainly governed by *Treponema* species. Of the 20 different phylotypes identified from DD biopsies, *Treponema phagedenis*-like, *T. medium*/*T. vincentii*-like and *T. denticola*/*T. pedis*-like phylotypes appears to be highly associated with the disease progression [1,2].

Specific niches of infection reservoirs and transmission routes, however still remains to be clarified in order to target and control the spread of the infection.

To address the issue of infection reservoirs of DD, we surveyed for evidence of treponemes in the content of the bovine gastrointestinal tract (feces), in DD lesions, and in the dairy farm environment (slurry).

## Conclusion

We surveyed for the presence of DD *Treponema* in environmental samples and individual scrapings from (DD- and mammary) lesions. We were not able to identify DD *Treponema* in the environmental samples (field, slurry and individual feces samples).

The phylogenetic compositions of the environmental samples were remarkably alike and significantly different from the DD-associated *Treponema*, figure 2.

The DD lesions and the ulcerative mammary dermatitis sample proved to be much more diverse and dominated by species previously identified in DD biopsies [1,2].

The question remains if this is a sensitivity issue – or if the reservoir of these bacteria is actually to be found elsewhere in the dairy herd environment.


## Methods

Samples from potential reservoirs were collected and subsequently DNA was purified by QIAamp®DNA stool mini kit. The hyper-variable regions V3 and V4 of 16S rRNA were PCR amplified as described in [1], and subsequently sequenced on a MiSeq, Illumina. Sequences were analyzed by BION-meta (<https://app.box.com/BION>)



**Figure 1. Healthy and digital dermatitis hoofs**

Severe lesions are easily identified (b,c) in comparison to a healthy hoof (a). Typical DD lesions characterized by irregular, raw raised areas that often develop keratinaceous hairlike projections (b, c).



**References:** 1. Klitgaard K, Boye M, Capion N, Jensen TK: Evidence of multiple *Treponema* phylotypes involved in bovine digital dermatitis as shown by 16S rRNA gene analysis and fluorescence in situ hybridization. *J Clin Microbiol* 2008, **46**(9):3012-3020.  
2. Evans NJ, Timofte D, Isherwood DR, Brown JM, Williams JM, Sherlock K, Lehane MJ, Murray RD, Birtles RJ, Hart CA *et al*: Host and environmental reservoirs of infection for bovine digital dermatitis treponemes. *Vet Microbiol* 2012, **156**(1-2):102-109.

**Acknowledgements:** Anastasia Isbrand is acknowledged for her technical support in the laboratory. Furthermore we would like to thank The Danish Cattle Federation for financial support of the project.

